

A note on the pressure of strong solutions to the Stokes system in bounded and exterior domains

Jörg Wolf
Department of Mathematics
Humboldt University Berlin
joerg@math.hu-berlin.de

Abstract

We consider the Stokes problem in an exterior domain $\Omega \subset \mathbb{R}^n$ with an external force $\mathbf{f} \in L^s(0, T; \mathbf{W}^{k, r}(\Omega))$ ($k \in \mathbb{N}, 1 < r < \infty$). In the present paper we show that in contrast to \mathbf{u} the boundary regularity of the pressure can be improved according to the differentiability of \mathbf{f} up to order k . In particular, this implies that the pressure is smooth with respect to $x \in \Omega$ if \mathbf{f} is smooth with respect to $x \in \Omega$.

Keywords Stokes equations, exterior domain, boundary regularity

Mathematics subject classification 35Q30, 76D03.

1 Introduction

Let $\Omega \subset \mathbb{R}^n$ ($n \in \mathbb{N}, n \geq 2$) be an exterior domain, i. e. $\mathbb{R}^n \setminus \overline{\Omega}$ is a bounded domain in \mathbb{R}^n . Let $0 < T < +\infty$. Set $Q = \Omega \times (0, T)$. In the present paper we consider the Stokes problem

$$(1.1) \quad \operatorname{div} \mathbf{u} = 0 \quad \text{in } Q$$

$$(1.2) \quad \partial_t \mathbf{u} - \Delta \mathbf{u} = -\nabla p + \mathbf{f} \quad \text{in } Q,$$

$$(1.3) \quad \mathbf{u} = 0 \quad \text{on } \partial\Omega \times (0, T), \quad \lim_{|x| \rightarrow \infty} \mathbf{u}(x, \cdot) = 0,$$

$$(1.4) \quad \mathbf{u}(0) = 0 \quad \text{in } \Omega,$$

where $\mathbf{u} = (u^1, \dots, u^n)$ denotes the unknown velocity of the fluid, p the unknown pressure and \mathbf{f} the given external force. The Stokes problem has been extensively studied in the past. In particular, for the case Ω is the half space or an C^2 domain with compact boundary the L^p -theory is well-known. Based on potential theory in [14] Solonnikov proved that for every $\mathbf{f} \in \mathbf{L}^q(Q)$ there exists a unique solution (\mathbf{u}, p) to (1.1)–(1.4) such that $\partial_t \mathbf{u}, \nabla^2 \mathbf{u} \in \mathbf{L}^q(Q)$, and $\nabla p \in \mathbf{L}^p(Q)$. By using the semi group approach, similar results have been obtained in [5], [6], [3]. For the corresponding estimates on the pressure we refer to [13]. An optimal result for the anisotropic case when \mathbf{f} belongs to $L^s(0, T; \mathbf{L}^q(\Omega))$ has been proved in [7] for the cases $\Omega = \mathbb{R}^n$, $\Omega = \mathbb{R}_+^n$, and a C^2 domain Ω with compact boundary.

By standard arguments from the regularity theory of parabolic equations one gets the regularity \mathbf{u} and p in dependence of the regularity of the right-hand side \mathbf{f} in time and space. However, if \mathbf{f} is only smooth in $x \in \Omega$ it is not clear whether \mathbf{u} is smooth in x up to the boundary. In the present paper we will see that such a property at least holds for the pressure p , which

is due to the fact that $\Delta p = 0$ if $\operatorname{div} \mathbf{f} = 0$. More precisely, the condition $\mathbf{f} \in L^s(0, T; \mathbf{W}^{k, q}(\Omega))$ ($1 < s, q, < +\infty; k \in \mathbb{N}$) implies $\nabla p \in L^s(0, T; \mathbf{W}^{k, q}(\Omega))$. Note that our result relies essentially on the fact that the initial data is zero. In general our result may not be true as there is a counter-example obtained in [9]. More precisely, there exists an initial data, and a solution \mathbf{u}, p to the Stokes system such that $\|\nabla \mathbf{u}(t)\|_{\mathbf{L}^2}$ is continuous as $t \rightarrow 0^+$, while the corresponding estimate on the pressure $\|p(t)\|_{L^2}$ may blow up as $t \rightarrow 0^+$.

First we shall introduce the basic notations regarding the function spaces used throughout the paper. By $W^{k, q}(\Omega), W_0^{k, q}(\Omega)$ we denote the usual Sobolev spaces. Vector functions and spaces of vector valued functions will be denoted by bold face letters, i. e. we write $\mathbf{L}^q(\Omega), \mathbf{W}^{k, q}(\Omega)$, etc. instead of $L^q(\Omega; \mathbb{R}^n), W^{k, q}(\Omega; \mathbb{R}^n)$, etc. In addition, we use the following spaces of solenoidal functions

$$\begin{aligned}\mathbf{L}_\sigma^q(\Omega) &= \text{closure of } \mathcal{C}_{0, \sigma}^\infty(\Omega) \text{ w.r.t. the norm } \|\cdot\|_{L^q} \\ \mathbf{W}_{0, \sigma}^{k, q}(\Omega) &= \text{closure of } \mathcal{C}_{0, \sigma}^\infty(\Omega) \text{ w.r.t. the norm } \|\cdot\|_{W^{k, q}},\end{aligned}$$

where $\mathcal{C}_{0, \sigma}^\infty(\Omega)$ stands for the space of all smooth solenoidal vector fields with compact support in Ω . Given a Banach space X by $L^q(0, T; X)$ we denote the space of Bochner measurable functions $f : (0, T) \rightarrow X$ such that

$$\begin{aligned}\|f\|_{L^q(0, T; X)}^q &= \int_0^T \|f(t)\|_X^q dt < +\infty \quad \text{if } 1 \leq q < +\infty, \\ \|f\|_{L^\infty(0, T; X)} &= \operatorname{ess\,sup}_{t \in (0, T)} \|f(t)\|_X < +\infty \quad \text{if } q = +\infty.\end{aligned}$$

Now, let us introduce the notion of a strong solution to (1.1)–(1.4).

Definition 1.1 Let $\mathbf{f} \in L^s(0, T; \mathbf{L}^q(\Omega))$ ($1 < s, q < +\infty$). A pair (\mathbf{u}, p) is called a *strong solution* to (1.1)–(1.4) if $\mathbf{u} \in L^s(0, T; \mathbf{W}_{0, \sigma}^{1, q}(\Omega)), p \in L^s(0, T; L_{\operatorname{loc}}^1(\overline{\Omega}))$ and

$$\partial_i \partial_j \mathbf{u}, \partial_t \mathbf{u}, \nabla p \in L^s(0, T; \mathbf{L}^q(\Omega)), \quad i, j = 1, 2, 3,$$

such that (1.1), (1.2) holds a. e. in Q , while (1.4) is fulfilled such that $\mathbf{u} = 0$ a. e. in $\Omega \times \{0\}$.

For the existence of a strong solution to (1.1)–(1.4) cf. in [7].

Our main result is the following

Theorem 1 Let $\Omega \subset \mathbb{R}^3$ be an exterior domain or a bounded domain with $\partial\Omega \in C^{2+k}$ ($k \in \mathbb{N}$). For $\mathbf{f} \in L^s(0, T; \mathbf{W}^{k, q}(\Omega))$ ($1 < s, q < +\infty; k \in \mathbb{N}$), let (\mathbf{u}, p) be the strong solution to (1.1)–(1.4). Then,

$$\nabla p \in L^s(0, T; \mathbf{W}^{k, q}(\Omega)).$$

In addition, there holds

$$(1.5) \quad \|\nabla^{k+1} p\|_{L^s(0, T; \mathbf{L}^q(\Omega))} \leq c \|\mathbf{f}\|_{L^s(0, T; \mathbf{W}^{k, q}(\Omega))},$$

where $c = \operatorname{const} > 0$ depending only on s, q, k and the geometric properties of $\partial\Omega$.

2 Remarks on the equation $\operatorname{div} \mathbf{v} = f$

Let $G \subset \mathbb{R}^n$ be a bounded domain, star-shaped with respect to a ball B_R . It is well known that for all $f \in L^q(G)$ with $(f)_G = 0$ ¹⁾ the equation $\operatorname{div} \mathbf{v} = f$ has a solution $\mathbf{v} \in \mathbf{W}_0^{1,q}(G)$ such that

$$\|\nabla \mathbf{v}\|_{L^q(G)} \leq c \|f\|_{L^q(G)}$$

with $c = \text{const} > 0$, depending on n, q and G (cf. [2], [8]). In fact, the constant c depends on the geometric property of G , namely the ratio of G which is defined by

$$\text{ratio}(G) := \frac{R_a(G)}{R_i(G)},$$

where

$$\begin{aligned} R_a(G) &= \inf\{R > 0 \mid \exists B_R(x_0) : G \subset B_R(x_0)\}, \\ R_i(G) &= \sup\{r > 0 \mid \exists B_r(x_0) : G \text{ is star-shaped w.r.t } B_r(x_0)\}. \end{aligned}$$

For instance $\text{ratio}(G) = 1$ if G is a ball, and $\text{ratio}(G) = \sqrt{n}$ if G is a cube. Moreover, the ratio is invariant under translation and scaling, i. e.

$$\text{ratio}(\lambda G) = \text{ratio}(G) \quad \forall \lambda > 0.$$

Now, let G such that $2 < R_i(G) < 3$. In particular, G is star shaped with respect to a ball $B_2 = B_2(x_0)$. Without loss of generality we may assume that $x_0 = 0$. Let $\phi \in C_0^\infty(B_2)$. We define

$$\mathcal{B}_\phi f(x) = \int_{\mathbb{R}^n} f(x-y) \mathbf{K}_\phi(x, y) dy, \quad x \in \mathbb{R}^n, \quad f \in C_0^\infty(G),$$

where

$$\mathbf{K}_\phi(x, y) = \frac{y}{|y|^n} \int_0^\infty \phi\left(x + r \frac{y}{|y|^n}\right) (|y| + r)^{n-1} dr, \quad (x, y) \in \mathbb{R}^n \times (\mathbb{R}^n \setminus \{0\}).$$

As in [2], [8] it has been proved that $\mathcal{B}_\phi f \in C_0^\infty(G)$ for all $f \in C_0^\infty(G)$. In addition, there holds

$$(2.1) \quad \|\nabla^k \mathcal{B}_\phi f\|_{L^q(G)} \leq c \|\nabla^{k-1} f\|_{L^q(G)} \quad \forall f \in C_0^\infty(G)$$

with a constant depending on n, k, q, ϕ and $\text{ratio}(G)$ only. Furthermore, there holds

$$(2.2) \quad \operatorname{div} \mathcal{B}_\phi f = f \int_{B_1} \phi(y) dy - \phi \int_G f(y) dy \quad \text{in } G.$$

In particular, if $\int_{B_1} \phi(y) dy = 1$ and $\int_G f(y) dy = 0$ then $\mathbf{v} = \mathcal{B}_\phi f$ solves the equation $\operatorname{div} \mathbf{v} = f$.

Finally, by (2.1) we may extend \mathcal{B}_ϕ to an operator $\mathcal{L}(W^{k-1,q}(G), \mathbf{W}^{k,q}(G))$ denoted again by \mathcal{B}_ϕ .

¹⁾ Let $A \subset \mathbb{R}^n$ be a measurable set with $\text{mes}(A) > 0$. Given $v \in L^1(A)$ by $(v)_A$ we denote the mean value $\frac{1}{\text{mes}(A)} \int_A v(x) dx$.

Let $i, j \in \{1, \dots, n\}$. Observing, that

$$\begin{aligned}\partial_j \mathcal{B}_\phi(\partial_i f) &= \partial_i \partial_j \mathcal{B}_\phi(f) - \partial_j \mathcal{B}_{\partial_i \phi}(f) \quad \text{in } G, \\ \partial_i \partial_j \mathcal{B}_\phi(f) &= \partial_i \mathcal{B}_\phi(\partial_j f) + \partial_i \mathcal{B}_{\partial_j \phi}(f) \quad \text{in } G,\end{aligned}$$

we see that

$$\partial_j \mathcal{B}_\phi(\partial_i f) = \partial_i \mathcal{B}_\phi(\partial_j f) + \partial_i \mathcal{B}_{\partial_j \phi}(f) - \partial_j \mathcal{B}_{\partial_i \phi}(f) \quad \text{in } G.$$

By the aid of (2.1), and Poincaré's inequality, using the above identity, we get

$$(2.3) \quad \|\partial_j \mathcal{B}_\phi(\partial_i f)\|_{L^q(G)} \leq c(\|\partial_j f\|_{L^q(G)} + \|f\|_{L^q(G)}) \quad \forall f \in W_0^{1,q}(G),$$

$$(2.4) \quad \begin{cases} \|\nabla^2 \partial_j \mathcal{B}_\phi(\partial_i f)\|_{L^q(G)} \leq c(\|\partial_i \nabla_* \nabla f\|_{L^q(G)} + \|\partial_j \partial_n \partial_n f\|_{L^q(G)} + \|\nabla^2 f\|_{L^q(G)})^2 \\ \forall f \in W_0^{3,q}(G), \end{cases}$$

where $c = \text{const} > 0$, depending on n, q and $\text{ratio}(G)$.

Now, let G be a bounded domain, star-shaped with respect to a ball B . Let $R := \frac{1}{2}R_i(G)$. Thus, there exist $B_R(x_0)$ such that G is star shaped to the ball $B_R(x_0)$. Without loss of generality we may assume that $x_0 = 0$. Let $\phi \in C_0^\infty(B_1)$ with $\int_{B_1} \phi(y) dy = 1$. We define

$\mathcal{B} : W_0^{k-1,q}(G) \rightarrow W_0^{k,q}(G)$ by setting

$$\mathcal{B}(f)(x) = R \mathcal{B}_\phi(\tilde{f})\left(\frac{x}{R}\right), \quad x \in G, \quad f \in W_0^{k-1,q}(G),$$

where $\tilde{f}(y) = f(Ry)$ ($y \in R^{-1}G$). Using the transformation formula of the Lebesgue integral, in view of (2.1), we see that

$$(2.5) \quad \begin{aligned} \|\nabla^k \mathcal{B}(f)\|_{L^q(G)} &= R^{n/q-k+1} \|\nabla^k \mathcal{B}_\phi(\tilde{f})\|_{L^q(R^{-1}G)} \leq c R^{n/q-k+1} \|\nabla^{k-1} \tilde{f}\|_{L^q(R^{-1}G)} \\ &= c \|\nabla^{k-1} f\|_{L^q(G)}, \end{aligned}$$

where $c = \text{const} > 0$ depends on n, q and $\text{ratio}(R^{-1}G) = \text{ratio}(G)$. In addition, from (2.3), and (2.4) we deduce

$$(2.6) \quad \|\partial_j \mathcal{B}(\partial_i f)\|_{L^q(G)} \leq c(\|\partial_j f\|_{L^q(G)} + R^{-1} \|f\|_{L^q(G)}) \quad \forall f \in W_0^{1,q}(G),$$

$$(2.7) \quad \begin{cases} \|\nabla^2 \partial_j \mathcal{B}(\partial_i f)\|_{L^q(G)} \leq c(\|\partial_i \nabla_* \nabla f\|_{L^q(G)} + \|\partial_j \partial_n \partial_n f\|_{L^q(G)} + R^{-1} \|\nabla^2 f\|_{L^q(G)}) \\ \forall f \in W_0^{3,q}(G), \end{cases}$$

($i, j = 1, \dots, n$) with a constant c , depending on n, q and $\text{ratio}(G)$ only. Furthermore, from (2.2) we get

$$(2.8) \quad \text{div } \mathcal{B}(f)(x) = f(x) - \phi\left(\frac{x}{R}\right) R^{-n} \int_G f(y) dy \quad \text{for a.e. } x \in G.$$

²⁾ Here ∇_* denotes the reduced gradient $(\partial_1, \dots, \partial_{n-1})$.

3 Proof of Theorem 1

Proof 1° By decomposing the right-hand side into a solenoidal field, and a gradient field, we are able to reduce the problem to the case $\operatorname{div} \mathbf{f} = 0$. Let $\mathbf{E} : \mathbf{W}^{k,q}(\Omega) \rightarrow \mathbf{W}^{k,q}(\mathbb{R}^n)$ denote an extension operator such that

$$\|\mathbf{E}\mathbf{v}\|_{\mathbf{W}^{k,q}(\mathbb{R}^n)} \leq c\|\mathbf{v}\|_{\mathbf{W}^{k,q}(\Omega)} \quad \forall \mathbf{v} \in \mathbf{W}^{k,q}(\Omega).$$

Let $\mathbf{P} : \mathbf{W}^{k,q}(\mathbb{R}^n) \rightarrow \mathbf{W}_{0,\sigma}^{k,q}(\mathbb{R}^n)$ denote the Helmholtz-Leray projection. Given $\mathbf{v} \in \mathbf{W}^{k,q}(\Omega)$ we have

$$\mathbf{v} = \mathbf{P}\mathbf{E}\mathbf{v} + (I - \mathbf{P})\mathbf{E}\mathbf{v} \quad \text{a.e. in } \Omega.$$

In addition, there exists a constant $c > 0$ depending only on n, q, k and Ω such that

$$(3.1) \quad \|\mathbf{P}\mathbf{E}\mathbf{v}\|_{\mathbf{W}^{k,q}(\Omega)} + \|(I - \mathbf{P})\mathbf{E}\mathbf{v}\|_{\mathbf{W}^{k,q}(\Omega)} \leq c\|\mathbf{v}\|_{\mathbf{W}^{k,q}(\Omega)} \quad \forall \mathbf{v} \in \mathbf{W}^{k,q}(\Omega).$$

Now, for $\mathbf{f} \in L^s(0, T; \mathbf{W}^{k,r}(\Omega))$ let (\mathbf{u}, p) be a strong solution to (1.1)–(1.4). Observing $I - \mathbf{P} = \nabla(\Delta^{-1} \operatorname{div})$ recalling the definition of \mathbf{E} we get

$$\mathbf{E}\mathbf{f} = \mathbf{P}\mathbf{E}\mathbf{f} + (I - \mathbf{P})\mathbf{E}\mathbf{f} = \mathbf{P}\mathbf{E}\mathbf{f} + \nabla(\Delta^{-1} \operatorname{div} \mathbf{E}\mathbf{f}) \quad \text{a.e. in } Q.$$

Since, $\nabla(\Delta^{-1} \operatorname{div} \mathbf{E}\mathbf{f}) = (\Delta^{-1} \nabla \operatorname{div} \mathbf{E}\mathbf{f}) \in L^s(0, T; \mathbf{W}^{k,q}(\mathbb{R}^n))$ we see that $\mathbf{P}\mathbf{E}\mathbf{f} \in L^s(0, T; \mathbf{W}^{k,q}(\mathbb{R}^n))$. Thus, we can replace \mathbf{f} by the restriction of $\mathbf{P}\mathbf{E}\mathbf{f}$ on Q , and p by the restriction of $-\Delta^{-1} \operatorname{div} \mathbf{E}\mathbf{f} + p$ on Q . Hence, in what follows without loss of generality we may assume that

$$(3.2) \quad \operatorname{div} \mathbf{f} = 0, \quad \text{and} \quad \Delta p = 0 \quad \text{a.e. in } Q.$$

2° Secondly, we recall a well-known result by Giga and Sohr [7] which is the following

Lemma 3.1 *Let $\Omega = \mathbb{R}^n$, $\Omega = \mathbb{R}_+^n$, Ω bounded or Ω an exterior domain with $\partial\Omega \in C^2$. For every $\mathbf{g} \in L^s(0, T; \mathbf{L}_\sigma^q(\Omega))$ ($1 < s, q, < +\infty$) there exists a unique solution $(\mathbf{v}, \pi) \in L^s(0, T; \mathbf{W}_{\text{loc}}^{2,q}(\Omega)) \times L^s(0, T; W_{\text{loc}}^{1,q}(\Omega))$ to the Stokes problem*

$$\begin{aligned} \partial_t \mathbf{v} - \Delta \mathbf{v} &= -\nabla \pi + \mathbf{g} \quad \text{and} \quad \operatorname{div} \mathbf{v} = 0 \quad \text{in } \Omega \times (0, T), \\ \mathbf{v} &= 0 \quad \text{on } \partial\Omega \times (0, T), \\ \mathbf{v}(0) &= 0 \quad \text{on } \Omega \times \{0\}, \end{aligned}$$

such that $\partial_t \mathbf{v}, \partial_i \partial_j \mathbf{v}, \nabla \pi \in L^s(0, T; \mathbf{L}^q(\Omega))$ ($i, j = 1, \dots, n$), and there holds,

$$(3.3) \quad \|\partial_t \mathbf{v}\|_{L^s(0,T;\mathbf{L}^q(\Omega))} + \|\nabla^2 \mathbf{v}\|_{L^s(0,T;\mathbf{L}^q(\Omega))} + \|\nabla \pi\|_{L^s(0,T;\mathbf{L}^q(\Omega))} \leq c\|\mathbf{g}\|_{L^s(0,T;\mathbf{L}^q(\Omega))},$$

where the constant c depends only on n, s, q and Ω .

As a consequence of Lemma 3.1 we get the existence of a unique solution $(\mathbf{u}, p) \in L^s(0, T; \mathbf{W}_{\text{loc}}^{2,q}(\Omega)) \times L^s(0, T; W_{\text{loc}}^{1,q}(\Omega))$ to the Stokes system (1.1)–(1.4), such that

$$(3.4) \quad \|\partial_t \mathbf{u}\|_{L^s(0,T;\mathbf{L}^q(\Omega))} + \|\nabla^2 \mathbf{u}\|_{L^s(0,T;\mathbf{L}^q(\Omega))} + \|\nabla p\|_{L^s(0,T;\mathbf{L}^q(\Omega))} \leq c\|\mathbf{f}\|_{L^s(0,T;\mathbf{L}^q(\Omega))}.$$

3° *Local estimates* We restrict ourself to case that Ω is an exterior domain. The opposite case can be treated in a similar way. Clearly, $G := \mathbb{R}^n \setminus \overline{\Omega}$ is a bounded domain. Let G', G'' are

bounded open sets such that $\overline{G} \subset G'$ and $\overline{G'} \subset G''$. Set $\Omega'' = \mathbb{R}^3 \setminus \overline{G''}$ and $\Omega' = \mathbb{R}^3 \setminus \overline{G'}$. Then, let $\zeta \in C^\infty(\mathbb{R}^3)$ denote a cut-off function such that $\zeta \equiv 1$ on Ω'' , and $\zeta \equiv 0$ in G' . In particular, $\text{supp}(\nabla \zeta) \subset G'' \setminus G'$. Observing $\text{div}(\mathbf{u}(t)\zeta) = \mathbf{u}(t) \cdot \nabla \zeta$, it follows that $\text{supp}(\mathbf{u}(t) \cdot \nabla \zeta) \subset \subset G'' \setminus G'$ for a. e. $t \in (0, T)$.

Next, let $1 < R < +\infty$ such that $G'' \subset B_R$. By $\mathcal{B} : W_0^{k-1,q}(B_R) \rightarrow \mathbf{W}_0^{k,q}(B_R)$ we denote the Bogowskii operator defined in Section 2. We now define

$$\mathbf{z}(t) = \mathcal{B}(\mathbf{u}(t) \cdot \nabla \zeta), \quad t \in [0, T].$$

Let $t \in (0, T)$. Since $\int_{B_R} \mathbf{u}(t) \cdot \nabla \zeta dx = 0$, in view of (2.8) we have

$$\text{div } \mathbf{z}(t) = \mathbf{u}(t) \cdot \nabla \zeta \quad \text{a. e. in } B_R.$$

Thanks to (2.6), recalling that $\text{ratio}(B_R) = 1$, there exists a constant $c > 0$ depending only on q and n such that

$$\|\mathbf{z}(t)\|_{\mathbf{W}^{3,q}(B_R)} \leq c \|\mathbf{u}(t) \cdot \nabla \zeta\|_{W^{2,q}(B_R)} \quad \text{for a. e. } t \in (0, T).$$

Making use of the embedding $\mathbf{W}_0^{3,q}(B_R) \hookrightarrow \mathbf{W}^{3,q}(\mathbb{R}^n)$ the above inequality implies that $\mathbf{z} \in L^s(0, T; \mathbf{W}^{3,q}(\mathbb{R}^n))$. Together with (3.4), and the Sobolev-Poincaré inequality we obtain

$$(3.5) \quad \|\mathbf{z}\|_{L^s(0,T;\mathbf{W}^{3,q}(\mathbb{R}^n))} \leq c \|\mathbf{u}\|_{L^s(0,T;\mathbf{W}^{2,q}(\Omega \cap B_R))} \leq c \|\mathbf{f}\|_{L^s(0,T;\mathbf{L}^q(\Omega))}.$$

By an analogous reasoning taking into account $\partial_t \mathbf{z} = \mathcal{B}(\partial_t \mathbf{u} \cdot \nabla \zeta)$ a. e. in $\mathbb{R}^n \times (0, T)$ we see that $\partial_t \mathbf{z} \in L^s(0, T; \mathbf{W}^{1,q}(\mathbb{R}^n))$. In addition, by virtue of (3.4) we obtain

$$(3.6) \quad \|\partial_t \mathbf{z}\|_{L^s(0,T;\mathbf{W}^{1,q}(\mathbb{R}^3))} \leq c \|\partial_t \mathbf{u}\|_{L^s(0,T;\mathbf{L}^q(\Omega))} \leq c \|\mathbf{f}\|_{L^s(0,T;\mathbf{L}^q(\Omega))}.$$

Next, let $k \in \{1, \dots, n\}$ be fixed. We define

$$\begin{cases} \mathbf{v}(x, t) = \partial_k(\mathbf{u}(x, t)\zeta(x) - \mathbf{z}(x, t)), & (x, t) \in (G'' \setminus G') \times (0, T), \\ \mathbf{v}(x, t) = -\partial_k \mathbf{z}(x, t), & (x, t) \in (\mathbb{R}^n \setminus (G'' \setminus G')) \times (0, T), \end{cases}$$

and

$$\begin{cases} \pi(x, t) = \partial_k(p(x, t)\zeta(x)), & (x, t) \in (G'' \setminus G') \times (0, T), \\ \pi(x, t) = 0, & (x, t) \in (\mathbb{R}^n \setminus (G'' \setminus G')) \times (0, T). \end{cases}$$

Then the pair (\mathbf{v}, π) solves the Stokes system

$$\begin{aligned} \text{div } \mathbf{v} &= 0 & \text{in } \mathbb{R}^n \times (0, T), \\ \partial_t \mathbf{v} - \Delta \mathbf{v} &= -\nabla \pi + \mathbf{g} & \text{in } \mathbb{R}^n \times (0, T), \\ \mathbf{v} &= 0 & \text{on } \mathbb{R}^n \times \{0\}, \end{aligned}$$

where

$$\begin{aligned} \mathbf{g} &= (p - p_{B_R})\nabla \zeta - 2\partial_k(\nabla \mathbf{u} \cdot \nabla \zeta) - \partial_k(\mathbf{u}\Delta \zeta) \\ &\quad - \partial_k \partial_t \mathbf{z} + \partial_k \Delta \mathbf{z} + \partial_k(\mathbf{f}\zeta) \quad \text{a. e. in } \mathbb{R}^n \times (0, T). \end{aligned}$$

In view of (3.3), (3.5), and (3.6) we see that $\mathbf{g} \in L^s(0, T; \mathbf{L}^q(\mathbb{R}^n))$. In addition, there holds

$$\|\mathbf{g}\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}^n))} \leq c \|\mathbf{f}\|_{L^s(0, T; \mathbf{W}^{1, q}(\Omega))}.$$

Thus, applying Lemma 3.1 with $\Omega = \mathbb{R}^n$, and using the last inequality we see that

$$\begin{aligned} & \|\partial_t \mathbf{v}\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}^n))} + \|\nabla^2 \mathbf{v}\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}^n))} + \|\nabla \pi\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}^n))} \\ & \leq c \|\mathbf{g}\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}^n))} \\ & \leq c \|\mathbf{f}\|_{L^s(0, T; \mathbf{W}^{1, q}(\Omega))}. \end{aligned}$$

Recalling the definition of \mathbf{v} , making use of (3.5), (3.6), and (3.4), we infer from above

$$\begin{aligned} & \|\zeta \partial_t \partial_k \mathbf{u}\|_{L^s(0, T; \mathbf{L}^q(\Omega))} + \|\zeta \nabla^2 \partial_k \mathbf{u}\|_{L^s(0, T; \mathbf{L}^q(\Omega))} + \|\zeta \nabla \partial_k p\|_{L^s(0, T; \mathbf{L}^q(\Omega))} \\ & \leq c \|\mathbf{f}\|_{L^s(0, T; \mathbf{W}^{1, q}(\Omega))}. \end{aligned}$$

Iterating the above argument k times, we get

$$\begin{aligned} & \|\partial_t \mathbf{u}\|_{L^s(0, T; \mathbf{W}^{k, q}(\Omega'))} + \|\mathbf{u}\|_{L^s(0, T; \mathbf{W}^{k+2, q}(\Omega'))} + \|\nabla p\|_{L^s(0, T; \mathbf{W}^{k, q}(\Omega'))} \\ (3.7) \quad & \leq c \|\mathbf{f}\|_{L^s(0, T; \mathbf{W}^{k, q}(\Omega))} \end{aligned}$$

($k \in \mathbb{N}$), where $c = \text{const} > 0$, depending on s, q, k , and Ω only.

4° *Boundary regularity* Let $x_0 \in \partial\Omega$. Up to translation and rotation we may assume that $x_0 = 0$ and $\mathbf{n}(0) = -\mathbf{e}_n$, where $\mathbf{n}(0)$ denotes the outward unite normal on Ω at x_0 . According to our assumption on the boundary of Ω there exists $0 < R < +\infty$, and $h \in C^{2+k}(B'_R)$ such that

- (i) $\partial\Omega \cap (B'_R \times (-R, R)) = \{(y', h(y')); y' \in B'_R\}$;
- (ii) $\{(y', y_n); y' \in B'_R, h(y') < y_n < h(y') + R\} \subset \Omega$;
- (iii) $\{(y', y_n); y' \in B'_R, -R + h(y') < y_n < h(y')\} \subset \Omega^c$ ³⁾.

Set $U_R = B'_R \times (-R, R)$, $U_R^+ = B'_R \times (0, R)$, and define $\Phi : U_R \rightarrow \Phi(U_R)$ by

$$\Phi(y) = (y', h(y') + y_n)^\top, \quad y \in U_R.$$

Elementary,

$$\begin{aligned} D\Phi(y) &= \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1 & 0 \\ \partial_1 h(y) & \partial_2 h(y) & \dots & \partial_{n-1} h(y) & 1 \end{pmatrix}, \\ (D\Phi(y))^{-1} &= \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1 & 0 \\ -\partial_1 h(y) & -\partial_2 h(y) & \dots & -\partial_{n-1} h(y) & 1 \end{pmatrix}. \end{aligned}$$

³⁾ Here $y' = (y_1, \dots, y_{n-1}) \in \mathbb{R}^{n-1}$, and B'_R denotes the two dimensional ball $\{(y_1, \dots, y_{n-1}) : y_1^2 + \dots + y_{n-1}^2 < R^2\}$.

For the outward unit normal at $x = \Phi(y)$ we have

$$\mathbf{n}(x) = \mathbf{N}(y) = \frac{(\partial_1 h(y), \dots, \partial_{n-1} h(y), -1)}{\sqrt{1 + |\nabla h(y)|^2}}, \quad y \in B'_R \times \{0\}.$$

In addition, one calculates

$$(3.8) \quad \partial_{x_i} \circ \Phi = \partial_{y_i} - (\partial_{x_i} h) \partial_{y_n} \quad \text{in } U_R, \quad i = 1, \dots, n^4).$$

We set $\mathbf{U} = \mathbf{u} \circ \Phi$, $P = p \circ \Phi$ and $\mathbf{F} = \mathbf{f} \circ \Phi$ a. e. in $U_R^+ \times (0, T)$. By the aid of (3.8) we easily get

$$(3.9) \quad (\operatorname{div}_x \mathbf{u}) \circ \Phi = \operatorname{div}_y \mathbf{U} - \nabla h \cdot \partial_{y_n} \mathbf{U} = 0,$$

$$(3.10) \quad (\Delta_x \mathbf{u}) \circ \Phi = \Delta_y \mathbf{U} - 2\nabla h \cdot \nabla_y \partial_{y_n} \mathbf{U} + |\nabla h|^2 \partial_{y_n} \partial_{y_n} \mathbf{U} - (\Delta h) \partial_{y_n} \mathbf{U},$$

$$(3.11) \quad (\nabla_x p) \circ \Phi = \nabla_y P - (\nabla h) \partial_{y_n} P,$$

a. e. in $U_R^+ \times (0, T)$. Firstly, owing to (3.9) from the equation (1.1) we get

$$(3.12) \quad \operatorname{div}_y \mathbf{U} = \nabla h \cdot \partial_{y_n} \mathbf{U} \quad \text{a. e. in } U_R^+ \times (0, T),$$

and with help of (3.10) and (3.11) the equation (1.2) turns into

$$(3.13) \quad \begin{aligned} \partial_t \mathbf{U} - \Delta \mathbf{U} = & -\nabla P + (\partial_{y_n} P) \nabla h - 2\nabla h \cdot \nabla \partial_{y_n} \mathbf{U} + |\nabla h|^2 \partial_{y_n} \partial_{y_n} \mathbf{U} \\ & - (\Delta h) \partial_{y_n} \mathbf{U} + \mathbf{F} \end{aligned}$$

a. e. in $U_R^+ \times (0, T)$.

Note that the assumption $\mathbf{n}(0) = -\mathbf{e}_n$ implies $\nabla h(0) = 0$. We now choose $0 < \delta < +\infty$ sufficiently small, which will be specified later. Since $\nabla h \in \mathbf{C}^0(U_R)$, there exists $0 < \rho < \frac{R}{2}$ such that

$$(3.14) \quad |\nabla h(y)| \leq \delta \quad \forall y \in U_{2\rho}.$$

Let $\zeta \in C_0^\infty(U_{2\rho})$ denote a cut-off function such that $0 \leq \zeta \leq 1$ in $U_{2\rho}$, and $\zeta \equiv 1$ on U_ρ . We define $\tilde{\mathbf{U}} : \mathbb{R}_+^n \times (0, T) \rightarrow \mathbb{R}^n$ by

$$\tilde{\mathbf{U}}(y, t) = \zeta(y) \mathbf{U}(y, t), \quad y \in U_{2\rho}^+ \times (0, T), \quad \tilde{\mathbf{U}}(y, t) = 0 \quad \text{if } y \in \mathbb{R}_+^n \setminus U_{2\rho}^+ \times (0, T).$$

Let $\mathcal{B} : W_0^{k-1, q}(U_{2\rho}^+) \rightarrow \mathbf{W}^{k, q}(\mathbb{R}_+^n)$ denote the Bogowskiĭ operator defined in Section 2. We set

$$\begin{aligned} \mathbf{z}_1(y, t) &= \mathcal{B}(\zeta \nabla h \cdot \partial_{y_n} \mathbf{U})(y, t), \\ \mathbf{z}_2(y, t) &= \mathcal{B}(\nabla \zeta \cdot \mathbf{U})(y, t), \quad (y, t) \in \mathbb{R}_+^n \times (0, T). \end{aligned}$$

Let $k \in \{1, \dots, n-1\}$ be fixed. We define

$$\begin{aligned} \mathbf{V}(y, t) &= \partial_k(\tilde{\mathbf{U}}(y, t) - \mathbf{z}_1(y, t) - \mathbf{z}_2(y, t)), \\ \Pi(y, t) &= \partial_k(\zeta(y) P(y, t)), \end{aligned}$$

⁴⁾ Since h is independent on y_n there holds $\partial_{x_n} \circ \Phi = \partial_{y_n}$.

$(y, t) \in \mathbb{R}_+^n \times (0, T)$. Observing that

$$\int_{U_{2\rho}^+} \zeta \nabla h \cdot \partial_n \mathbf{U}(t) + \nabla \zeta \cdot \mathbf{U}(t) dy = \int_{U_{2\rho}^+} \operatorname{div}_y \tilde{\mathbf{U}}(t) dy = 0 \quad \text{for a. e. } t \in (0, T),$$

by the aid of (2.8) we calculate

$$(3.15) \quad \operatorname{div}_y \mathbf{V} = \partial_k \left(\zeta \nabla h \cdot \partial_n \mathbf{U} + \nabla \zeta \cdot \mathbf{U} - \zeta \nabla h \cdot \partial_n \mathbf{U} - \nabla \zeta \cdot \mathbf{U} \right) = 0$$

a. e. in $\mathbb{R}_+^n \times (0, T)$. In addition, taking into account (3.13), we find

$$\begin{aligned} \partial_t \mathbf{V} - \Delta \mathbf{V} &= \partial_k \left(\zeta \partial_t \mathbf{U} - \zeta \Delta \mathbf{U} - 2 \nabla \zeta \cdot \nabla \mathbf{U} - (\Delta \zeta) \mathbf{U} \right) \\ &\quad - \partial_k (\partial_t \mathbf{z}_1 - \Delta \mathbf{z}_1) - \partial_k (\partial_t \mathbf{z}_2 - \Delta \mathbf{z}_2) \\ &= -\nabla \Pi + \partial_k ((P - P_{U_{2\rho}^+}) \nabla \zeta) - \partial_k (2 \nabla \zeta \cdot \nabla \mathbf{U} + (\Delta \zeta) \mathbf{U}) \\ &\quad - \partial_k (\partial_t \mathbf{z}_1 - \Delta \mathbf{z}_1) - \partial_k (\partial_t \mathbf{z}_2 - \Delta \mathbf{z}_2) \\ &\quad + \partial_k \left(\zeta (\partial_n P) \nabla h - 2 \zeta \nabla h \cdot \nabla \partial_n \mathbf{U} + \zeta |\nabla h|^2 \partial_n \partial_n \mathbf{U} \right. \\ &\quad \left. - \zeta (\Delta h) \partial_n \mathbf{U} + \zeta \mathbf{F} \right). \end{aligned}$$

Thus, (\mathbf{V}, Π) solves the following Stokes system

$$\begin{aligned} \operatorname{div} \mathbf{V} &= 0 & \text{in } \mathbb{R}_+^n \times (0, T), \\ \partial_t \mathbf{V} - \Delta \mathbf{V} &= -\nabla \Pi + \mathbf{G} & \text{in } \mathbb{R}_+^n \times (0, T), \\ \mathbf{V} &= 0 & \text{on } \partial \mathbb{R}_+^n \times (0, T), \end{aligned}$$

where $\mathbf{G} = \mathbf{G}_1 + \dots + \mathbf{G}_6$ with

$$\begin{aligned} \mathbf{G}_1 &= \partial_{y_k} ((P - P_{U_{2\rho}^+}) \nabla \zeta), \\ \mathbf{G}_2 &= -\partial_k (2 \nabla \zeta \cdot \nabla \mathbf{U} + (\Delta \zeta) \mathbf{U}), \\ \mathbf{G}_3 &= -\partial_k (\partial_t \mathbf{z}_1 - \Delta \mathbf{z}_1), \\ \mathbf{G}_4 &= -\partial_k (\partial_t \mathbf{z}_2 - \Delta \mathbf{z}_2), \\ \mathbf{G}_5 &= \partial_k \left(\zeta (\partial_n P) \nabla h - 2 \zeta \nabla h \cdot \nabla \partial_n \mathbf{U} + \zeta |\nabla h|^2 \partial_n \partial_n \mathbf{U} \right), \\ \mathbf{G}_6 &= \partial_k (-\zeta (\Delta h) \partial_n \mathbf{U} + \zeta \mathbf{F}). \end{aligned}$$

In what follows we shall establish some important estimates of \mathbf{z}_1 and \mathbf{z}_2 , where we will make essential use of the properties of \mathcal{B} (cf. Section 2). Starting with \mathbf{z}_1 , we write $\mathbf{z}_1 = \mathbf{z}_{1,1} + \mathbf{z}_{1,2}$, where

$$\mathbf{z}_{1,1} = \mathcal{B}(\partial_n (\zeta \nabla h \cdot \mathbf{U})), \quad \mathbf{z}_{1,2} = -\mathcal{B}((\partial_n \zeta) \nabla h \cdot \mathbf{U}).$$

Let $t \in (0, T)$ be fixed. Using (2.5), (2.6) with $j = k, i = n$ and $f = \zeta \nabla h \cdot \mathbf{U}$, and observing $\partial_t \mathcal{B} = \mathcal{B} \partial_t$, we see that

$$\begin{aligned} \|\partial_t \partial_k \mathbf{z}_1(t)\|_{L^q(\mathbb{R}_+^n)} &\leq \|\partial_t \partial_k \mathbf{z}_{1,1}(t)\|_{L^q(\mathbb{R}_+^n)} + \|\partial_t \partial_k \mathbf{z}_{1,2}(t)\|_{L^q(\mathbb{R}_+^n)} \\ &\leq c \|\partial_t \partial_k (\zeta \nabla h \cdot \mathbf{U})(t)\|_{L^q(\mathbb{R}_+^n)} + c \rho^{-1} \|\partial_t \mathbf{U}(t)\|_{L^q(\mathbb{R}_+^n)} \\ &\leq c \delta \|\partial_t \partial_k \tilde{\mathbf{U}}(t)\|_{L^q(\mathbb{R}_+^n)} + c (\|h\|_{C^2} + \rho^{-1}) \|\partial_t \mathbf{U}(t)\|_{L^q(U_{2\rho}^+)}. \end{aligned}$$

Taking the above inequality to the s -th power, and integrating the resulting equation in time over $(0, T)$, we get

$$(3.16) \quad \begin{aligned} & \|\partial_t \partial_k \mathbf{z}_1\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}_+^n))} \\ & \leq c\delta \|\partial_t \partial_k \tilde{\mathbf{U}}\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}_+^n))} + c(\|h\|_{C^2} + \rho^{-1}) \|\partial_t \mathbf{U}\|_{L^s(0, T; \mathbf{L}^q(U_R^+))}. \end{aligned}$$

On the other hand, using (2.5), (2.7) with $j = k, i = n$, and $f = \zeta \nabla h \cdot \mathbf{U}(t)$, we see that

$$\begin{aligned} \|\nabla^2 \partial_k \mathbf{z}_1(t)\|_{\mathbf{L}^q(\mathbb{R}_+^n)} & \leq c \|\partial_n \nabla_* \nabla(\zeta \nabla h \cdot \mathbf{U})(t)\|_{\mathbf{L}^q(\mathbb{R}_+^n)} + c \|\partial_n \partial_n \partial_k(\zeta \nabla h \cdot \mathbf{U})(t)\|_{\mathbf{L}^q(\mathbb{R}_+^n)} \\ & \quad + c\rho^{-1} \|\nabla^2(\zeta \nabla h \cdot \mathbf{U})(t)\|_{\mathbf{L}^q(\mathbb{R}_+^n)} + c \|\nabla^2((\partial_n \zeta) \nabla h \cdot \mathbf{U})(t)\|_{\mathbf{L}^q(\mathbb{R}_+^n)}. \end{aligned}$$

By means of product rule and Poincaré's inequality we find

$$\|\nabla^2 \partial_k \mathbf{z}_1(t)\|_{\mathbf{L}^q(\mathbb{R}_+^n)} \leq c\delta \|\nabla^2 \nabla_* \tilde{\mathbf{U}}(t)\|_{\mathbf{L}^q(\mathbb{R}_+^n)} + c(\|h\|_{C^3} + \rho^{-1}) \|\nabla^2 \mathbf{U}(t)\|_{\mathbf{L}^q(U_R^+)}.$$

We now take the above inequality to the s -th power, integrating the result in time over $(0, T)$, we obtain

$$(3.17) \quad \begin{aligned} \|\nabla^2 \partial_k \mathbf{z}_1\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}_+^n))} & \leq c\delta \|\nabla^2 \nabla_* \tilde{\mathbf{U}}\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}_+^n))} \\ & \quad + c(\|h\|_{C^3} + \rho^{-1}) \|\nabla^2 \mathbf{U}\|_{L^s(0, T; \mathbf{L}^q(U_R^+))}. \end{aligned}$$

By an analogous reasoning, making use of (2.5), and Poincaré's inequality, we infer

$$(3.18) \quad \begin{aligned} & \|\partial_t \mathbf{z}_2\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}_+^n))} + \|\nabla^2 \mathbf{z}_2\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}_+^n))} \\ & \leq c\rho^{-1} \left(\|\partial_t \mathbf{U}\|_{L^s(0, T; \mathbf{L}^q(U_R^+))} + \|\nabla^2 \mathbf{U}\|_{L^s(0, T; \mathbf{L}^q(U_R^+))} \right). \end{aligned}$$

We are now in a position to estimate $\mathbf{G}_1, \dots, \mathbf{G}_6$. First by virtue of Poincaré's inequality we easily estimate

$$\|\mathbf{G}_1\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}_+^n))} \leq c\rho^{-1} \|\nabla P\|_{L^s(0, T; \mathbf{L}^q(U_R^+))}.$$

Analogously,

$$\|\mathbf{G}_2\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}_+^n))} \leq c\rho^{-1} \|\nabla^2 \mathbf{U}\|_{L^s(0, T; \mathbf{L}^q(U_R^+))}.$$

Next, with the help of (3.16), (3.17), and (3.18) we see that

$$\begin{aligned} & \|\mathbf{G}_3\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}_+^n))} + \|\mathbf{G}_4\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}_+^n))} \\ & \leq c\delta \left(\|\partial_t \partial_k \tilde{\mathbf{U}}\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}_+^n))} + \|\nabla^2 \nabla_* \tilde{\mathbf{U}}\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}_+^n))} \right) \\ & \quad + c(\|h\|_{C^3} + \rho^{-1}) \left(\|\partial_t \partial_k \tilde{\mathbf{U}}\|_{L^s(0, T; \mathbf{L}^q(U_R^+))} + \|\nabla^2 \nabla_* \tilde{\mathbf{U}}\|_{L^s(0, T; \mathbf{L}^q(U_R^+))} \right). \end{aligned}$$

Then applying the product rule, and using Poincaré's inequality, we get

$$\begin{aligned} \|\mathbf{G}_5\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}_+^n))} & \leq c\delta \left(\|\nabla \Pi\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}_+^n))} + \|\nabla_* \nabla^2 \tilde{\mathbf{U}}\|_{L^s(0, T; \mathbf{L}^q(\mathbb{R}_+^n))} \right) \\ & \quad + c(\|h\|_{C^2} + \rho^{-1}) \left(\|\nabla P\|_{L^s(0, T; \mathbf{L}^q(U_R^+))} + \|\nabla^2 \mathbf{U}\|_{L^s(0, T; \mathbf{L}^q(U_R^+))} \right). \end{aligned}$$

Finally, we estimate

$$\begin{aligned} \|\mathbf{G}_6\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^n))} &\leq c(\|h\|_{C^3} + \rho^{-1}) \left(\|\nabla P\|_{L^s(0,T;\mathbf{L}^q(U_R^+))} + \|\nabla^2 \mathbf{U}\|_{L^s(0,T;\mathbf{L}^q(U_R^+))} \right) \\ &\quad + c\rho^{-1} \|\mathbf{F}\|_{L^s(0,T;\mathbf{L}^q(U_R^+))} + c\|\partial_k \mathbf{F}\|_{L^s(0,T;\mathbf{L}^q(U_R^+))}. \end{aligned}$$

Appealing to Lemma 3.1 (cf. [7]) for the case $\Omega = \mathbb{R}_+^n$ using the above estimates for $\mathbf{G}_1, \dots, \mathbf{G}_6$, we obtain

$$\begin{aligned} &\|\partial_t \mathbf{V}\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^3))} + \|\nabla^2 \mathbf{V}\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^3))} + \|\nabla \Pi\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^3))} \\ &\leq c\|\mathbf{G}_1 + \dots + \mathbf{G}_6\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^n))} \\ &\leq c\delta \left(\|\partial_t \nabla_* \tilde{\mathbf{U}}\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^3))} + \|\nabla^2 \nabla_* \tilde{\mathbf{U}}\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^3))} + \|\nabla \Pi\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^3))} \right) \\ &\quad + c(\|h\|_{C^3} + \rho^{-1}) \left(\|\partial_t \mathbf{U}\|_{L^s(0,T;\mathbf{L}^q(U_R^+))} + \|\nabla^2 \mathbf{U}\|_{L^s(0,T;\mathbf{L}^q(U_R^+))} \right. \\ &\quad \left. + \|\nabla P\|_{L^s(0,T;\mathbf{L}^q(U_R^+))} + \|\mathbf{F}\|_{L^s(0,T;\mathbf{W}^{1,q}(U_R^+))} \right). \end{aligned}$$

Recalling $\mathbf{V} = \partial_k(\tilde{\mathbf{U}} - \mathbf{z}_1 - \mathbf{z}_2)$, making use of (3.16), (3.17) and (3.18), from the last inequality we infer

$$\begin{aligned} &\|\partial_t \nabla_* \tilde{\mathbf{U}}\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^3))} + \|\nabla^2 \nabla_* \tilde{\mathbf{U}}\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^3))} + \|\nabla \Pi\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^3))} \\ &\leq c_0 \delta \left(\|\partial_t \nabla_* \tilde{\mathbf{U}}\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^3))} + \|\nabla^2 \nabla_* \tilde{\mathbf{U}}\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^3))} + \|\nabla \Pi\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^3))} \right) \\ &\quad + c_1 \left(\|\partial_t \mathbf{U}\|_{L^s(0,T;\mathbf{L}^q(U_R^+))} + \|\nabla^2 \mathbf{U}\|_{L^s(0,T;\mathbf{L}^q(U_R^+))} \right. \\ (3.19) \quad &\quad \left. + \|\nabla P\|_{L^s(0,T;\mathbf{L}^q(U_R^+))} + \|\mathbf{F}\|_{L^s(0,T;\mathbf{W}^{1,q}(U_R^+))} \right), \end{aligned}$$

where $c_0 = c_0(n, q, s)$ and $c_1 = c_1(n, q, s, \|h\|_{C^3}, \rho)$. On the other hand, recalling the definition of \mathbf{U}, P , and \mathbf{F} , with the help of (3.10), (3.11), and (3.7) we find

$$\begin{aligned} &\|\partial_t \mathbf{U}\|_{L^s(0,T;\mathbf{L}^q(U_R^+))} + \|\nabla^2 \mathbf{U}\|_{L^s(0,T;\mathbf{L}^q(U_R^+))} \\ (3.20) \quad &\quad + c\|\nabla P\|_{L^s(0,T;\mathbf{L}^q(U_R^+))} + \|\mathbf{F}\|_{L^s(0,T;\mathbf{W}^{1,q}(U_R^+))} \leq c\|\mathbf{f}\|_{L^s(0,T;\mathbf{W}^{1,q}(\Omega))} \end{aligned}$$

with a constant c depending on n, q, s and h . Now, in (3.19) we take $\delta = \frac{1}{2c_0}$ and estimate the right-hand side of (3.19) by the aid of (3.20). This leads to

$$\begin{aligned} &\|\partial_t \nabla_* \tilde{\mathbf{U}}\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^3))} + \|\nabla^2 \nabla_* \tilde{\mathbf{U}}\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^3))} + \|\nabla \Pi\|_{L^s(0,T;\mathbf{L}^q(\mathbb{R}_+^3))} \\ &\leq c_2 \|\mathbf{f}\|_{L^s(0,T;\mathbf{W}^{1,q}(\Omega))}, \end{aligned}$$

where $c_2 = c_2(n, q, s, \|h\|_{C^3}, \rho)$.

By a standard iteration argument we obtain

$$\begin{aligned} &\|\partial_t \nabla_*^k \mathbf{U}\|_{L^s(0,T;\mathbf{L}^q(U_\rho^+))} + \|\nabla^2 \nabla_*^k \mathbf{U}\|_{L^s(0,T;\mathbf{L}^q(U_\rho^+))} + \|\nabla \nabla_*^k P\|_{L^s(0,T;\mathbf{L}^q(U_\rho^+))} \\ (3.21) \quad &\leq c\|\mathbf{f}\|_{L^s(0,T;\mathbf{W}^{1,q}(\Omega))}, \end{aligned}$$

where $c = \text{const}$ depending only on $n, q, s, k, \|h\|_{C^{k+2}}$ and ρ .

5° *Estimation of the full pressure gradient* Recalling that $\Delta_x p = 0$, with the help of (3.10) we calculate

$$\begin{aligned} 0 = \Delta_x p \circ \Phi &= \Delta_y P - 2\nabla h \cdot \nabla \partial_{y_n} P + |\nabla h|^2 \partial_{y_n} \partial_{y_n} P - (\Delta h) \partial_{y_n} P \\ &= (1 + |\nabla h|^2) \partial_n \partial_{y_n} P + \Delta'_y P - 2\nabla h \cdot \nabla_* \partial_n P - (\Delta h) \partial_{y_n} P^5 \end{aligned}$$

a. e. in U_R^+ . Thus,

$$(1 + |\nabla h|^2) \partial_{y_n} \partial_{y_n} P = -\Delta'_y P + 2\nabla h \cdot \nabla_* \partial_{y_n} P + (\Delta h) \partial_{y_n} P$$

a. e. in U_R^+ . From this identity along with (3.21) with $k = 1$ it follows that

$$\begin{aligned} \|\nabla_y^2 P\|_{L^s(0,T;\mathbf{L}^q(U_\rho^+))} &\leq c \left(\|\nabla_* \partial_{y_n} P\|_{L^s(0,T;\mathbf{L}^q(U_\rho^+))} + \|\nabla_y P\|_{L^s(0,T;\mathbf{L}^q(U_\rho^+))} \right) \\ &\leq c \|\mathbf{f}\|_{L^s(0,T;\mathbf{W}^{1,q}(\Omega))}. \end{aligned}$$

Choosing $\rho \in \left(0, \frac{R}{2}\right)$ sufficiently small, and applying the above argument k -times, we get

$$(3.22) \quad \|\nabla_y^{k+1} P\|_{L^s(0,T;\mathbf{L}^q(U_\rho^+))} \leq c \|\mathbf{f}\|_{L^s(0,T;\mathbf{W}^{k,q}(\Omega))}$$

with a constant c depending on $n, q, s, k, \|h\|_{C^{k+2}}$, and ρ .

Finally a standard covering argument, together with (3.22), and (3.7) gives the estimate (1.5), which completes the proof of the Theorem 1. \blacksquare

Acknowledgements The present research has been supported by the German Research Foundation (DFG) through the project WO1988/1-1; 612414.

References

- [1] Adams, R. A., **Sobolev spaces**, Academic Press, Boston 1978.
- [2] M. E. Bogowskii, *Solution to the First Boundary Value Problem for the Equation of Continuity of an Incompressible Medium*, Soviet Math. Dokl. **20** (1979), 1094–1098.
- [3] W. Borchers, H. Sohr, *On the semigroup of the Stokes operator for exterior domains*, Math. Z. **196** (1987), 415–425.
- [4] Farwig, R.; Kozono, H.; Sohr, H., *An L^q -approach to Stokes and Navier-Stokes equations in general domains*, Acta Math. **195**, 21–53 (2005).
- [5] Y. Giga, *Analyticity of the semi group generated by the Stokes operator in L_r spaces*, Math. Z. **178** (1981), 297–329.
- [6] Y. Giga, *Domains of fractional powers of the Stokes operator in L_r spaces*, Arch. Rational Mech. Anal. **89** (1985), 251–265.
- [7] Y. Giga, H. Sohr, *Abstract L^p estimates for the Cauchy problem with applications to the Navier-Stokes equations in exterior domains*, J. Funct. Anal. **102** (1991), 72–94.
- [8] P. G. Galdi, **An Introduction to the Mathematical Theory of the Navier-Stokes Equations, Linearized Steady Problems**, Vol. 38, Springer, New York, 1994.

⁵⁾ Here Δ'_y stands for the differential operator $\partial_{y_1} \partial_{y_1} + \dots + \partial_{y_{n-1}} \partial_{y_{n-1}}$.

- [9] J. G. Heywood, O. D. Walsh, **A counter-example concerning the pressure in the Navier-Stokes equation, as $t \rightarrow 0^+$** , Pacific J. Math. **164**, No. 2, 1994.
- [10] Hopf, E., *Über die Anfangswertaufgabe für die Hydrodynamischen Grundgleichungen*, Math. Nachr. **4**, 213 (1950/1951).
- [11] Leray, J., *Sur le mouvements d'un liquide visqueux emplissant l'espace*, Acta Math. **63**, 193-284 (1934).
- [12] H. Sohr, **The Navier-Stokes Equations. An Elementary Functional Analytic Approach**, Birkhäuser Advanced Texts, Birkhäuser Verlag, Basel, 2001. 1994.
- [13] H. Sohr, W. von Wahl, *On the regularity of the pressure of weak solutions of Navier-Stokes equations* Arch. Math. **46** (1986), 428–439.
- [14] V. A. Solonnikov, *Estimates for solutions of non-stationary Navier-Stokes equations*, J. Soviet Math **8** (1977), 467–523.